




ИНСТИТУТ ЯДЕРНОЙ ФИЗИКИ СО АН СССР

25

V.G. Dubrovsky, B.G. Konopelchenko

ON THE INTERRELATION
BETWEEN THE SOLUTIONS OF
THE MKP AND KP EQUATIONS
VIA MIURA TRANSFORMATION

PREPRINT 91-30



НОВОСИБИРСК

On the Interrelation
between the Solutions of
the mKP and KP Equations
via Miura Transformations

V.G. Dubrovsky, B.G. Konopelchenko

Institute of Nuclear Physics
630090, Novosibirsk, USSR

ABSTRACT

It is shown that the classes of exact solutions with functional parameters and rational solutions of the modified Kadomtsev—Petviashvili (mKP) equation and the Kadomtsev—Petviashvili (KP) equation are connected by the 2+1-dimensional Miura transformation. The correspondence between more particular classes of solutions of the mKP and the KP equations via Miura transformation is established.

BETWEEN THE SOLUTIONS OF
THE mKP AND KP EQUATIONS
VIA MIURA TRANSFORMATION

PREPRINT 81-03

1. INTRODUCTION

The Miura transformation [1] between the modified Korteweg-de Vries (mKdV) and the Korteweg-de Vries (KdV) equations has played an important role both in the discovery the inverse scattering transform (IST) method [2] and in the further understanding of the properties of these equations (see e.g. [3—5]). It reveals a deep interrelation between the algebraic properties of the mKdV and KdV equations and their hierarchies [3—5].

The 2+1-dimensional integrable generalizations of the KdV and mKdV equations are given by the well-known Kadomtsev—Petviashvili (KP) equation

$$U_t + U_{xxx} + 6UU_x + 3\sigma^2 \partial_x^{-1} U_{yy} = 0 \quad (1)$$

and the modified KP (mKP) equation

$$V_t + V_{xxx} - 3\sigma^2 \left(\frac{1}{2} V^2 V_x - V_x \partial_x^{-1} V_y + \partial_x^{-1} V_{yy} \right) = 0 \quad (2)$$

which has been introduced within the different approaches in [6, 7]. Here $\sigma^2 = \pm 1$. The mKP and KP equations are connected by the 2+1-dimensional generalization of the Miura transformation. Namely, if the function V obeys the mKP equation (2) then the function

$$U = -\frac{1}{2} \sigma^2 \partial_x^{-1} V_y - \frac{1}{2} \sigma V_x - \frac{1}{4} \sigma^2 V^2 \quad (3)$$

obeys the KP equation [6, 7]. Similar to the 1+1-dimensional case the Miura transformation (3) deeply interrelates the algebraic structures associated with the mKP and KP equations [8].

In the present paper the properties of the classes of exact solutions of the mKP and the KP equations under the Miura map (3) are studied. We will show that the solutions of the mKP equation (2) with functional parameters are converted under the map (3) into the solutions of the KP equation (1) with, essentially, the same functional parameters. The Miura map (3) transforms the rational solutions of the mKP equation into the rational solutions of the KP equation. The interrelation between the plane solitons, decreasing and plane lumps, nonsingular and singular, real and complex solutions of the mKP and KP equations are established. Both cases $\sigma^2 = -1$ and $\sigma^2 = +1$ are considered.

2. GENERAL FORMULAE

The classes of exact solutions of the KP equation, including solutions with functional parameters, plane solitons, singular rational solutions and lumps, are well-known (see e.g. [3, 5]). For the mKP equation similar classes of exact solutions has been constructed recently in [9]. All the details about the properties of the solutions of the KP and mKP equations can be found in these papers.

The correspondence between the solutions of the mKP and KP equations can be established directly in the terms of V and U with the use of the map (3). But it is more convenient and transparent to do this using the mKP eigenfunction χ . The first linear problem for the mKP equation (2) is of the form [9]

$$\sigma\Psi_y + \Psi_{xx} + \sigma V\Psi_x = 0. \quad (4)$$

The eigenfunction $\chi(x, y, t; \lambda)$ relevant for the formulation and solving the inverse spectral problem for the mKP equation is introduced via [9]

$$\Psi \stackrel{\text{def}}{=} \chi \exp\left(\frac{ix}{\lambda} + \frac{y}{\sigma\lambda^2}\right). \quad (5)$$

It obeys the equation

$$\sigma\chi_y + \chi_{xx} + \frac{2i}{\lambda}\chi_x + \frac{i\sigma V}{\lambda}\chi + \sigma V\chi_x = 0. \quad (6)$$

The function $\chi(x, y, t; \lambda)$ can be canonically normalized ($\chi \xrightarrow{\lambda \rightarrow \infty} 1$) and the reconstruction formula for the potential V is of the form [9]

$$V(x, y, t) = -\frac{2}{\sigma} \frac{\partial}{\partial x} \ln \chi_0 \quad (7)$$

where χ_0 is the first of coefficients of the Taylor series expansion of χ near origin:

$$\chi(x, y, t; \lambda) = \chi_0(x, y, t) + \lambda\chi_1(x, y, t) + \lambda^2\chi_2(x, y, t) + \dots \quad (8)$$

The equations of the inverse problem for (6) have the wide classes of exact solutions which give rise via (7) to the classes of exact solutions of the mKP equations [9].

The formula (7) is very convenient for the analysis of the Miura map (3). Substituting (7) into (3), one easily gets

$$U = -\frac{\sigma(\chi_0^{-1})_y + (\chi_0^{-1})_{xx}}{\chi_0^{-1}} \quad (9)$$

that clearly demonstrates that U is, indeed, the solution of the KP equation. Moreover using the relation

$$\sigma\chi_{0y} + \chi_{0xx} + 2i\chi_{1x} + i\sigma V\chi_1 + \sigma V\chi_{0x} = 0 \quad (10)$$

which arises from the substitution of expansion (8) into (6), one can transform the r.h.s. of (9) into the simpler form. Namely,

$$U = -2i \frac{\partial}{\partial x} \left(\frac{\chi_1}{\chi_0} \right). \quad (11)$$

This formula gives us a simple way for the calculating the solutions of the KP equation using the known functions χ_0 and χ_1 for the mKP equation.

It is easy to see that the Miura map (3) transforms the real-valued solutions of the mKP-I equation ($\sigma=i$) into the complex-valued solutions of the KP equation ($\sigma=i$) while in the case $\sigma=1$ the Miura transformation (3) connects the real-valued solutions of the mKP and KP equations. The Miura transformation (9) maps the

nonsingular solutions of the mKP equation into the nonsingular solutions of the KP equation.

3. CORRESPONDENCE BETWEEN THE SOLUTIONS WITH FUNCTIONAL PARAMETERS AND RATIONAL SOLUTIONS

The most general classes of exact solutions of the mKP equations include the solutions with functional parameters and rational in x, y, t solutions. Here we will consider the correspondence between them.

The solutions of the mKP equation with functional parameters are of the form [9]

$$V = -\frac{2}{\sigma} \frac{\partial}{\partial x} \ln(\det \tilde{A} A^{-1}) \quad (12)$$

where

$$\begin{aligned} A_{kl} &= \delta_{kl} - \frac{1}{2i} \partial_x^{-1} \xi_{kx} \eta_l, \\ \tilde{A}_{kl} &= \delta_{kl} + \frac{1}{2i} \partial_x^{-1} \xi_k \eta_{lx} \end{aligned} \quad (13)$$

and

$$\chi_0 = \det(\tilde{A} A^{-1}), \quad \chi_1 = \text{tr}(B A^{-1}) \quad (14)$$

where

$$B_{kl} = i \partial_x \tilde{A}_{kl} = \frac{1}{2} \xi_k \eta_{lx}. \quad (15)$$

Here $\xi_k(x, y, t)$ and $\eta_l(x, y, t)$ are the solutions of the linearized mKP equation

$$\xi_t + \xi_{xxx} + 3\sigma^2 \partial_x^{-1} \xi_{yy} = 0 \quad (16)$$

of the form

$$\begin{aligned} \xi_k(x, y, t) &= \iint_C d\lambda \wedge d\bar{\lambda} f_k(\lambda, \bar{\lambda}) \exp\left(\frac{ix}{\lambda} + \frac{y}{\sigma\lambda^2} + \frac{4it}{\lambda^3}\right), \\ \eta_l(x, y, t) &= \iint_C d\lambda \wedge d\bar{\lambda} g_k(\lambda, \bar{\lambda}) \exp\left(\frac{ix}{\lambda} + \frac{y}{\sigma\lambda^2} + \frac{4it}{\lambda^3}\right), \end{aligned} \quad (17)$$

where $f_k(\lambda, \bar{\lambda})$ and $g_k(\lambda, \bar{\lambda})$ are arbitrary complex functions. The integral ∂_x^{-1} in (13), (14) is defined in such a way that r.h.s. of (13) and (14) exist.

Now let us apply the Miura transformation to the solutions (12). The formula (11) gives

$$U = 2 \frac{\partial}{\partial x} \frac{\text{tr}(\tilde{A}_x A^{-1})}{\det(\tilde{A} A^{-1})}. \quad (18)$$

Note that the matrices $\tilde{A}_x A^{-1}$, $1 - A\tilde{A}^{-1} - \tilde{A}_x \tilde{A}^{-1}$ and $1 - A\tilde{A}^{-1}$ have the rank one. For the rank one matrices one has well-known identity

$$\det(1 + F) = 1 + \text{tr} F. \quad (19)$$

Using (19) and another well-known matrix identity

$$\frac{\partial}{\partial x} \ln \det F = \text{tr}(F_x F^{-1}) \quad (20)$$

one gets:

$$\begin{aligned} \frac{\text{tr}(\tilde{A}_x A^{-1})}{\det(\tilde{A} A^{-1})} &= \frac{\det(1 + \tilde{A}_x A^{-1}) - 1}{\det(\tilde{A} A^{-1})} = \\ &= \det(1 - (1 - A\tilde{A}^{-1} - \tilde{A}_x \tilde{A}^{-1})) - \det(1 - (1 - A\tilde{A}^{-1})) = \\ &= \text{tr}(\tilde{A}_x \tilde{A}^{-1}) = \frac{\partial}{\partial x} \ln \det \tilde{A}. \end{aligned} \quad (21)$$

So we finally obtain the solutions of the KP equation

$$U = 2 \frac{\partial^2}{\partial x^2} \ln \det \tilde{A} \quad (22)$$

with the functional parameters, where \tilde{A} is given by (13). The formula (22) after the identification

$$\xi_l^{\text{KP}} = \xi_l, \quad \eta_k^{\text{KP}} = \eta_{kx} \quad (23)$$

exactly coincides with the known formula for the solutions of the KP equation with functional parameters (see e.g. [3]).

The linear parts of the KP and the mKP equations coincide. So, the same set (up to the change (23)) of solutions of the linear equation (16) parametrize the classes of exact solutions (12) and (22) of the mKP and KP equations, respectively. The Miura map (3)

connects these classes of solutions without, in essence, change the parameters ξ_l and η_k .

Similar situation takes place for the rational solutions. The general rational solutions of the mKP equation are of the form [9].

$$V = -\frac{2}{\sigma} \frac{\partial}{\partial x} \ln \det(\tilde{A} A^{-1}) \quad (24)$$

where

$$A_{kl} = \delta_{kl} \left(x - \frac{2iy}{\sigma\lambda_k} + \frac{12t}{\lambda_k^2} + \gamma \right) + (1 - \delta_{kl}) \frac{i\lambda_l^2}{\lambda_k - \lambda_l},$$

$$\tilde{A}_{kl} = A_{kl} + i\lambda_l \quad (25)$$

and

$$\chi_0 = \det(\tilde{A} A^{-1}), \quad \chi_1 = i \operatorname{tr}(\mathcal{E} A^{-1}) \quad (26)$$

where $\mathcal{E}_{kl} = 1$ ($k, l = 1, \dots, N$).

The Miura map (3) convert the solutions (24) into the following solutions of the KP equation

$$U = 2 \frac{\partial}{\partial x} \frac{\operatorname{tr}(\mathcal{E} A^{-1})}{\det(\tilde{A} A^{-1})} \quad (27)$$

Taking into account that the matrices $1 - A\tilde{A}^{-1} - \mathcal{E}\tilde{A}^{-1}$ and $1 - A\tilde{A}^{-1}$ have rank one, and using the identities (19), (20), similar to the previous case, one gets

$$U = 2 \frac{\partial}{\partial x} \operatorname{tr}(\mathcal{E}\tilde{A}^{-1}). \quad (28)$$

Finally, using the properties of the matrix \tilde{A} (in particular, $(\tilde{A}_{pq})_x = \delta_{pq}$), one obtains

$$U = 2 \frac{\partial^2}{\partial x^2} \ln \det \tilde{A} \quad (29)$$

where \tilde{A} is given by (25), i.e.

$$\tilde{A}_{kl} = \delta_{kl} \left(x - \frac{2iy}{\sigma\lambda_k} + \frac{12t}{\lambda_k^2} + \gamma \right) + i(1 - \delta_{kl}) \frac{1}{\lambda_l^{-1} - \lambda_k^{-1}}. \quad (30)$$

The formulae (29), (30) coincide with the well-known formulae for the rational solutions of the KP equation (up to $\lambda_k \rightarrow \lambda_k^{-1}$) (see [3]).

4. CORRESPONDENCE BETWEEN PARTICULAR CLASSES OF SOLUTIONS. CASE $\sigma = i$.

Now we will consider more particular classes of solutions, including the real and nonsingular solutions.

4.1. We start with the case $\sigma = i$ ($\sigma^2 = -1$). The real nonsingular plane solitons of the mKP-I equation are given by the formula (12) with [9]

$$\xi_l = -2iR_l \exp(F(\lambda_l)), \quad \eta_k = -2i \exp(-F(\bar{\lambda}_k)) \quad (31)$$

where $\operatorname{Im} R_l = 0$ and $F(\lambda) \stackrel{\text{def}}{=} \frac{ix}{\lambda} - \frac{iy}{\lambda^2} + \frac{4it}{\lambda^3}$. It is not difficult to show using (22), that the real nonsingular plane solitons of the mKP-I equations are transformed into complex nonsingular plane solitons of the KP-I equation. In particular the simplest mKP-I soliton [9]

$$V = -\frac{8(\lambda_r/|\lambda|^2) \operatorname{sgn} R}{e^{2f} + (e^{-f} + (\lambda_r/\lambda) \operatorname{sgn} R) e^f} \quad (32)$$

is converted into the complex nonsingular soliton of the KP-I equation

$$U = \frac{8\lambda \lambda_r \operatorname{sgn} R}{|\lambda|^4 (e^{-f} + (\lambda/\lambda_r) e^f \operatorname{sgn} R)^2} \quad (33)$$

where

$$2f = ix(\lambda^{-1} - \bar{\lambda}^{-1}) - iy(\lambda^{-2} - \bar{\lambda}^{-2}) + 4it(\lambda^{-3} - \bar{\lambda}^{-3}) + \ln |R|. \quad (34)$$

The well-known plane real-valued nonsingular solitons of the KP-I are connected via Miura transformation (3) with the complex nonsingular plane solitons of the mKP-I equation. In particular the well-known one-soliton solution of the KP-I equation

$$U = \frac{2\lambda_r^2}{|\lambda|^4} \frac{1}{\operatorname{ch}^2 f} \quad (35)$$

is obtained from the complex nonsingular soliton

$$V = -\frac{4\lambda_r^2}{\lambda |\lambda|^2} \frac{1}{(e^{-f} + \frac{\bar{\lambda}}{\lambda} e^f) \operatorname{ch} f} \quad (36)$$

of the mKP-I equation, where f is given by (34).

4.2. The solutions of the mKP-I equation of the breather type, constructed in [9] are converted by the Miura transformation into the periodic in x or y solutions of the KP-I equation. In particular, the complex breather type solution of the mKP-I equation which can be obtained by the technique of the work [9] is of the form

$$V = 2i \frac{\partial}{\partial x} \ln \frac{\left(1 - \frac{e^f \sin \varphi}{\lambda_R}\right)^2 + \frac{e^{2f}}{\lambda_R^2 \lambda_I^2} (\lambda_R^2 + \lambda_I^2 \cos^2 \varphi)}{1 + \frac{2e^f \bar{\lambda}^2}{\lambda_R |\lambda|^2} \sin \varphi + \frac{\bar{\lambda}^2 e^{2f}}{\lambda_R^2 \lambda_I^2}} \quad (37)$$

where

$$f = -\frac{4\lambda_R \lambda_I}{|\lambda|^4} y + \ln |\lambda R|, \quad (38)$$

$$\varphi = \frac{2\lambda_R x}{|\lambda|^4} + \frac{8(\lambda_R^3 - 3\lambda_R \lambda_I^2) t}{|\lambda|^6} + \arg(R\lambda)$$

and R some complex constant is transformed into the following real solution of the KP-I equation

$$U = 2 \frac{\partial^2}{\partial x^2} \ln \left\{ \left(1 - \frac{e^f \sin \varphi}{\lambda_R}\right)^2 + \frac{e^{2f}}{\lambda_R^2 \lambda_I^2} (\lambda_R^2 + \lambda_I^2 \cos^2 \varphi) \right\}. \quad (39)$$

The solution (39) is the real-valued, nonsingular solution of the KP-I equation decreasing at $y \rightarrow \pm \infty$, and has a periodic wave character in x, t .

4.3. Another complex breather type solution of the mKP-I equation which can be obtained by the technique of the work [9] is of the form:

$$V = 2i \frac{\partial}{\partial x} \ln \frac{1 + ae^f \cos \varphi + \frac{(v_1 + v_2)^2}{16v_1 v_2} a^2 e^{2f}}{1 + \frac{a}{2} e^f \left(\frac{v_2}{v_1} e^{i\varphi} + \frac{v_1}{v_2} e^{-i\varphi} \right) + \frac{a^2}{4} e^{2f}} \quad (40)$$

where

$$f(x, t) = x \left(\frac{1}{v_1} - \frac{1}{v_2} \right) - 4t \left(\frac{1}{v_1^3} - \frac{1}{v_2^3} \right),$$

$$\varphi(y) = y \left(\frac{1}{v_1^2} - \frac{1}{v_2^2} \right)$$

and a some real constant, is converted by Miura transformation into

$$U = 2 \frac{\partial^2}{\partial x^2} \ln \left\{ 1 + ae^f \cos \varphi + \frac{a^2}{16} \frac{(v_1 + v_2)^2}{v_1 v_2} e^{2f} \right\}. \quad (41)$$

This real, nonsingular, periodic in y and soliton type in (x, t) solution of the KP-I equation has been found in [10].

5. THE CORRESPONDENCE BETWEEN THE LUMPS ($\sigma=i$)

The KP-I equation possesses the real decreasing lumps (see [3, 5]) while the mKP-I equation has both real decreasing lumps and real plane lumps [9].

5.1. The real decreasing lumps of the mKP-I equation are given by the formula (24) with [9]

$$N = 2n, \quad \lambda_{n+i} = \bar{\lambda}_i$$

$$\gamma_i = -\frac{i\lambda_i}{2} + c_i, \quad \gamma_{n+i} = -\frac{i\bar{\lambda}_i}{2} + \bar{c}_i \quad (42)$$

where $\lambda_i (i=1, \dots, n)$ are arbitrary isolated points outside the real axis and c_i are arbitrary constants.

It is not difficult to see that the corresponding rational solutions (29), (30) of the KP-I equation are complex and nonsingular. In particular, the real nonsingular decreasing lump of the mKP-I equation [9]

$$V = \frac{2\lambda \bar{X}^2 + 2\bar{\lambda} X^2 - \frac{|\lambda|^2 \lambda_R^3}{\lambda_I^2}}{\left(|X|^2 + \frac{|\lambda|^2 \lambda_R^2}{4\lambda_I^2}\right)^2 + \left(\frac{\lambda}{2} \bar{X} + \frac{\bar{\lambda}}{2} X\right)^2} \quad (43)$$

where

$$X = x - \frac{2y}{\lambda} + \frac{12t}{\lambda^2} + c,$$

$$c = c_R + ic_I$$

is converted into the complex nonsingular decreasing rational solution

$$U = \frac{|\lambda|^2 \lambda_R^2 - 2X^2 - 2\bar{X}^2 + 2\lambda_R^2 - 2i\bar{\lambda}\bar{X} - 2i\lambda X}{\lambda_j^2 \left[|X|^2 + \frac{|\lambda|^2 \lambda_R^2}{4\lambda_j^2} + i\left(\frac{\lambda}{2}\bar{X} + \frac{\bar{\lambda}}{2}X\right) \right]^2} \quad (44)$$

of the KP-I equation.

5.2. The real plane lumps of the mKP-I equation are given by (24), (25) where [9]

$$\text{Im } \lambda_i = 0, \quad \gamma_i = -\frac{i\lambda_i}{2} + c_i, \quad \text{Im } c_i = 0 \quad (i=1, \dots, N).$$

They are mapped by the Miura transformation (3) into the complex plane nonsingular rational solutions of the KP-I equation. For instance, the simplest plane lump of the mKP-I equation

$$V = \frac{2\alpha}{\left(x - \frac{2y}{\alpha} + \frac{12t}{\alpha^2} + x_0\right)^2 + \frac{\alpha^2}{4}} \quad (45)$$

where α is an arbitrary real constant is converted to the complex plane nonsingular lump:

$$U = 2 \frac{\frac{\alpha^2}{4} - \left(x - \frac{2y}{\alpha} + \frac{12t}{\alpha^2} + x_0\right)^2 + i\alpha \left(x - \frac{2y}{\alpha} + \frac{12t}{\alpha^2} + x_0\right)}{\left[\left(x - \frac{2y}{\alpha} + \frac{12t}{\alpha^2} + x_0\right)^2 + \frac{\alpha^2}{4}\right]^2} \quad (46)$$

5.3. At last one can show that the real decreasing lumps of the KP-I equations [3, 5] are obtained by the Miura transformation (3) from the complex rational nonsingular solutions of the mKP-I equation. In particular, the complex solution

$$V = 2i \left\{ \frac{X + \bar{X}}{|X|^2 + \frac{|\lambda|^4}{4\lambda_j^2}} - \frac{X + \bar{X} - 2i\lambda_R}{|X|^2 + \frac{|\lambda|^4}{4\lambda_j^2} - |\lambda|^2 - i(\lambda\bar{X} + \bar{\lambda}X)} \right\} \quad (47)$$

where

$$X = x - \frac{2y}{\lambda} + \frac{12t}{\lambda^2} + c$$

of the mKP-I equation is transformed into

$$U = \frac{|\lambda|^4 - 2X^2 - 2\bar{X}^2}{\lambda_j^2 \left(|X|^2 + \frac{|\lambda|^4}{4\lambda_j^2}\right)^2} \quad (48)$$

that is the well-known real decreasing lump of the KP-I equation [3, 5].

6. THE CASE $\sigma=1$

In this case the real valued solutions of the mKP-II equation are transformed by the Miura map into the real-valued solutions of the KP-II equation.

6.1. The real plane solitons of the mKP-II equation are given by the formulae (12), (13) with

$$\begin{aligned} \xi_i &= -2iR_i \exp(F(i\alpha_i)), \\ \eta_i &= -2\beta_i^{-1} \exp(-F(i\beta_i)) \end{aligned} \quad (49)$$

where R_i, α_i, β_i are arbitrary real constants. It is easy to see that the corresponding solutions of the KP-II equation are given by (22) with

$$\bar{A}_{mn} = \delta_{mn} - \frac{2R_n \exp[x(\alpha_n^{-1} - \beta_n^{-1}) - y(\alpha_n^{-2} - \beta_n^{-2}) - 4t(\alpha_n^{-3} - \beta_n^{-3})]}{\beta_n^2(\alpha_n^{-1} - \beta_n^{-1})} \quad (50)$$

that exactly coincides with the formula for the multi-soliton solutions of the KP-II equation (with $\alpha_n^{-1} \rightarrow \lambda_n$) [3]. In particular, the simplest real soliton of the mKP-II equation [9]

$$V = -\frac{2(\alpha - \beta)^2}{\alpha\beta^2} \frac{\varepsilon}{\left(e^{-t - \frac{\alpha}{\beta}\varepsilon t}\right) \left(e^{-t - \varepsilon t}\right)} \quad (51)$$

where

$$2f = (\alpha^{-1} - \beta^{-1})x - (\alpha^{-2} - \beta^{-2})y - 4(\alpha^{-3} - \beta^{-3})t + \ln \frac{2|R|}{|\beta - \alpha|},$$

$$\varepsilon = \text{sgn}\left(\frac{R}{\beta - \alpha}\right) \quad (52)$$

is converted into the well-known real plane soliton of the KP-II equation:

$$U = \frac{1}{2} \left(\frac{1}{\alpha} - \frac{1}{\beta} \right)^{-2} \text{ch}^{-2} \left(\frac{\tilde{f}}{2} \right) \quad (53)$$

where

$$\tilde{f} = (\alpha^{-1} - \beta^{-1})x - (\alpha^{-2} - \beta^{-2})y - 4(\alpha^{-3} - \beta^{-3})t + \ln \frac{2R\alpha}{\beta(\alpha - \beta)}. \quad (54)$$

The function (53) is nonsingular not only for those values of parameters $\alpha, \beta, \varepsilon$ ($\varepsilon < 0, \alpha/\beta > 0$) as the mKP-II plane soliton (51) but also for $\varepsilon > 0, \alpha/\beta < 0$ for which the soliton (51) is the singular one. The properties of the KP-II plane solitons are quite different in these two cases. Namely, the soliton (53) at $\varepsilon > 0, \alpha/\beta < 0$ (type I) possesses at $\alpha = -\beta$ the nontrivial 1+1-dimensional limit

$$U_{\text{KdV}} = \frac{2}{\alpha^2} \text{ch}^{-2} \frac{\varphi}{2}, \quad \varphi = \tilde{f} |_{\alpha = -\beta} \quad (55)$$

that is the standard KdV soliton while at $\varepsilon < 0, \alpha/\beta > 0$ (type II) the solution (53) has a trivial 1+1-dimensional limit $U|_{\alpha = \beta} = 0$.

So the Miura transformation (3) maps the bounded plane solitons of the mKP-II equation into the type II (pure 2+1-dimensional) plane solitons of the KP-II equation and the singular plane soliton of the mKP-II equation into the standard (type I) plane soliton of the KP-II equation.

This last property of the map (3) is similar to the property of the 1+1-dimensional Miura map $U = -\frac{1}{2}V_x - \frac{1}{4}V^2$ which, as it has been shown in [11], does not interrelates the rapidly decaying smooth solutions of the mKdV and KdV equations. This is quite clear from the consideration of the 1+1-dimensional limit of the 2+1-dimensional case. Indeed, the 1+1-dimensional limit of the solution (51) ($\alpha = -\beta$) looks like

$$V_{\text{mKdV}} = \frac{4\varepsilon}{\alpha} \text{sh}^{-1} 2\varphi, \quad \varphi = f |_{\alpha = -\beta} \quad (56)$$

that is the singular solution of the mKdV equation while the corresponding limit of the solution (53) ($\alpha = -\beta$) is given by (55). So the 1+1-dimensional Miura transformation maps the singular

solutions of the mKdV equation into the soliton of the KdV equation.

Similar situation takes place for general multi-soliton solutions of the mKP-II equations.

6.2. The rational solutions of the mKP-II equation are real-valued in the two cases [9]

- 1) $N = 2n, \quad \lambda_{k+n} = \bar{\lambda}_k, \quad \gamma_{k+n} = \bar{\gamma}_k \quad (k = 1, \dots, n);$
- 2) arbitrary $N, \quad \lambda_k = i\alpha_k \quad (\text{Im } \alpha_k = 0), \quad \gamma_k = \bar{\gamma}_k.$

But all these rational solutions of the mKP-II equation are singular.

They remain singular after the Miura transformation. In particular, the simplest singular plane lumps of the mKP-II equation

$$V = \frac{2\alpha}{\frac{\alpha^2}{4} - \left(x + \frac{2y}{\alpha} - \frac{12t}{\alpha^2} + x_0 \right)^2}, \quad (57)$$

where α is an arbitrary real constant is transformed into the solution

$$U = -\frac{2}{\left(x + \frac{2y}{\alpha} - \frac{12t}{\alpha^2} - \frac{\alpha}{2} + x_0 \right)^2}. \quad (58)$$

This the well-known singular solution of the KP-II equation [3, 5].

Note that the singularity's line $x = -\frac{2y}{\alpha} + \frac{12t}{\alpha^2} + \frac{\alpha}{2} - x_0$ of the solution (58) coincides with one of the singularity's line of the solution (57).

Similar situation takes place in the general case too. Comparing the formulae (24) and (29), we see that the singularities of the solutions of the KP-II equation are defined by the zeros of the det A and that these singularities present only the part of the singularities of the solution (24) of the mKP-II equation.

REFERENCES

1. R.M. Miura. J. Math. Phys., **9**, 1202 (1968).
2. C.S. Gardner, G.M. Green, M. Kruskal, R.M. Miura. Phys. Rev. Lett., **19**, 1095 (1967).
3. V.E. Zakharov, S.V. Manakov, S.P. Novikov, L.P. Pitaevski. Theory of Solitons.—Nauka, Moscow, 1980.
4. G.L. Lamb. Jr. Elements of Soliton Theory.—John Wiley & Sons, New York, 1980.
5. M.J. Ablowitz and H. Segur. Solitons and the Inverse Scattering Transform.—SIAM, Philadelphia, 1981.
6. B.G. Konopelchenko. Phys. Lett., **92A**, 323 (1982).
7. M. Jimbo, T. Miwa. Publ. RIMS, **19(3)**, 943 (1983).
8. B.G. Konopelchenko. Nonlinear Integrable Equations.—Lecture Notes in Physics, v.270, Springer, Berlin-Heidelberg, 1987.
9. B.G. Konopelchenko. V.G. Dubrovsky. Preprint INP 90-149 (1990); Stud. Appl. Math. (submitted).
10. V.E. Zakharov, S.V. Manakov. Funkt. Anal. Prilozh., **19(2)**, 11 (1985).
11. M.J. Ablowitz, M. Kruskal, H. Segur. J. Math. Phys., **20**, 999 (1979).

В.Г. Дубровский, Б.Г. Конопельченко

**О связи решений уравнений МКП и КП
преобразованием Миуры**

V.G. Dubrovsky, B.G. Konopelchenko

**On the Interrelation
between the Solutions of
the mKP and KP Equations
via Miura Transformations**

Ответственный за выпуск С.Г. Попов

Работа поступила 3 апреля 1991 г.
Подписано в печать 8 апреля 1991 г.
Формат бумаги 60×90 1/16 Объем 1,5 печ.л., 1,2 уч.-изд.л.
Тираж 250 экз. Бесплатно. Заказ № 30

*Набрано в автоматизированной системе на базе фото-
наборного автомата ФА1000 и ЭВМ «Электроника» и
отпечатано на ротапринтере Института ядерной физики
СО АН СССР,
Новосибирск, 630090, пр. академика Лаврентьева, 11.*